7th Grade Physics Pre-Comprehensive Exam Prep

The Long List of Things to Memorize

How to use this guide

- This is a list of items that must be memorized in order to have success on the precomprehensive exam in 7th grade physics.
- o You must be able to recite any of these definitions or equations instantaneously.
- For any item which you are not 100% confident that you have memorized, you should make a flashcard where the left hand column is the front side of the flashcard and the right hand column is the back of the flashcard.
- To effectively use flashcards, you must go over them every single day. This means 7 days a week from now until the pre-comprehensive examination, or until you are 100% confident that you have memorized the concept and will have it memorized forever.
 - o 10 minutes a day or less, EVERY DAY
- o To use flashcards: start with a single pile of cards. Go through them one at a time. Any that you do not get correct goes into a "review" pile, any that you do get correct goes into a "completed" pile. Once you have gone through the initial pile, pick up the review pile only and try it again. Continue to sort those into "complete" and "review" piles. Any word you get correct the second time goes into the "complete" pile, any you are still uncertain of goes into the "review" pile again. Go through the review pile 4 times.
- Any card which ends up in the "review" pile after 4 times is something you need to study. Read the front of the flashcard out loud, turn it over and read the back of it out loud. Repeat this process for a total of 5 times.
- o Repeat the entire process tomorrow.
- Any card which goes into the "complete" pile on the first time through the pile every day for a week should be removed from your pile of flashcards and you should consider it memorized.
- O We will not have class time to review vocabulary terms in class; you must spend time studying your vocabulary terms on your own.

Scientific Measurement and graphing

Qualitative measurement	o Answers questions other than "how many"
	 Describes an object
Quantitative measurement	o Answers the question "how many" or
	"how much"
	o Includes a number
Metric units of length	o meters (m)
Metric units of mass	o kilogram (kg)
Metric units of weight	o Newton (N)
Metric units of time	o second (s)
Metric units of volume	o meters cubed (m³) or liters (L)
Metric units of temperature	o Kelvin (K)
Metric prefix for 10 ⁹	o Giga (G-)
Metric prefix for 10 ⁶	o Mega (M-)
Metric prefix for 10 ³	o kilo (k-)
Metric prefix for 10 ²	o hecto (h-)
Metric prefix for 10 ¹	o deka (da-)
Metric prefix for 10 ⁻¹	o deci (d-)
Metric prefix for 10 ⁻²	o centi (c-)
Metric prefix for 10 ⁻³	o milli (m-)
Metric prefix for 10 ⁻⁶	o micro (μ-)
Metric prefix for 10 ⁻⁹	o nano (n-)
Equation for slope of a graph	$\circ m = \frac{rise}{run} = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$
Units for slope	o Units on the y axis divided by units on the
	x axis
How to make a graph	 Vertical axis is dependent variable
	 Horizontal axis is independent variable
	 Both axes need even scales

Position, Velocity, and Acceleration

definition of reference point	 a stationary point from which another object's position is measured typically used to determine if the second object is in motion
definition of position	 the vector quantity describing the location, both distance and direction, from a reference point to an object
definition of motion	 when the position of one object is changing when measured from a reference point

Definition of distance	how much ground was actually covered in a tripscalar
Definition of displacement	how far an object has moved from its' initial positionvector
definition of speed	o how fast an object is moving
units for speed	o m/s
Instantaneous speed	Speed at any one instant in timeFound using speedometer
Average speed	 Total distance travelled divided by total time that passed while travelling
Equation for average speed	$\begin{array}{c} \circ \ s = d/\Delta t \\ \circ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$
Definition of velocity	Change in displacement in a time intervalSpeed in a specific direction
Average Velocity Equation	$\circ v_{avg} = \frac{\Delta x}{total \ time}$
Constant velocity	\circ Acceleration is 0 m/s^2
	 Object has constant speed
	 Object does not change direction
Definition of acceleration	o Change in velocity
Positive Acceleration	Acceleration in the positive direction
Negative Acceleration	Acceleration in the negative direction
units for acceleration	$\circ m/s^2$
Equation with acceleration, initial velocity,	$\circ a = (v_f - v_i)/t$
final velocity, and time interval	$ \circ v_f = v_i + at$
Definition of resultant velocity	o The net total of velocities
velocity from a position time graph	 Velocity is the slope of a position time graph
Acceleration from a velocity time graph	 Acceleration is the slope of a velocity time graph

Forces and Newton's 1st and 2nd Laws of Motion

Statement of Newton's 1st law of motion	 An object in motion will remain in motion
	and an object at rest will remain at rest
	unless acted upon by an unbalanced force.
Definition of inertia	o A measurement of the mass of an object
Inertia determined by mass	 Only the amount of mass changes the
	amount of inertia
Equation for weight	$\circ F_{grav} = mg$
Definition of weight	The force of gravity on an object

GM_1m_2
$\circ F_g = \frac{GM_1m_2}{r^2}$
 inverse-square relationship of F_g and r
 G is the universal gravitational
constant (do not need to memorize
number)
 mass does not change with gravity
changes, but weight depends on the force
of gravity
 Drawing depicting all of the forces which
act on an object
$\circ \overrightarrow{F_{net}} = m\vec{a}_{net}$
 One value increases as another increases
o one value increases when another
decreases
o a push or a pull on an object
o Newton (N)
\circ 1N = 1kg*(1m/s ²)=1kgm/s ²
In order from strongest to weakest:
 Strong Nuclear Force
o Electromagnetism
Weak Nuclear force
o Gravity
 The result with all the (vector) forces that
act on an object are added together
o a force that occurs only when 2 objects are
in contact
o a force that occurs between 2 objects that
are not in contact
o quantities with a magnitude and a
direction
o quantities with only a magnitude
 When the net force on an object is 0N
whether it is moving or stationary
 2 types- dynamic and static
o when the forces add up to 0N
 When the forces do not add up to 0N
o force caused by a surface which is
perpendicular to the surface
o force caused by a rope
o force cause by surface which is parallel to
the surface
ο F _f =μF _N

Coefficient of friction	$\circ \mu$, the measure of how rough a surface is
	o It has no units
Cause of friction	o friction occurs because surfaces are not
	perfectly smooth

Newton's 3rd Law of Motion

Newton's 3rd law of motion	o every action (force) has a reaction
	(corresponding force) that is equal in
	magnitude and opposite in direction
Newton's 3 rd Law statement	o The force on object 2 by object 1 is equal
	in magnitude and opposite direction the
	force on object 2 by object 1.
properties of action and reaction forces	o they occur at exactly the same time
	o they are always the same size
	o they act between the same two objects
	o they each act on separate objects

Energy

Statement of the conservation of energy law	 Energy cannot be created nor destroyed, it can only be transformed from one type to another
Equation for PEgrav	⊙ PE _{grav} =mgh
Equation for KE	\circ KE = $\frac{1}{2}$ mv ²
Equation for Conservation of Mechanical Energy (isolated system)	\circ TME $_i$ = TME $_f$
Calculating total mechanical energy of a system	⊙TME = KE + PE
Definition of Energy	○ The ability to do work
Units for Energy	○ Joules (J)= 1kgm²/s²
Definition of Kinetic Energy	○ The energy of motion
Definition of Potential Energy	 Energy that is stored by an object due to its' position relative to another object or by a change in the object's shape
Definition of gravitational potential energy	 Energy an object possesses because of its position in a gravitational field
Definition of elastic potential energy	Energy of an object due to a change in the object's shape
Definition of chemical potential energy	 energy stored in an object due to relative positions of the molecules
Definition of electromagnetic energy	energy contained in an electromagnetic wave/photon

Definition of thermal energy	 energy due to the vibrations of atoms in an object
Sound Energy	 energy associated with the wave motion of particles
Definition of electric energy	 energy due to the relative positions or motion of electrically charged objects
Definition of nuclear energy	 energy contained within the nucleus; comes from bonding of protons and neutrons
Definition of work	o when a force causes a displacement
Equation for Work	$\circ W = Fd$
Units for work	o Joules (J) o Nm
Definition of Power	o the rate at which work is done by a force
Equation for power	$\circ P = \frac{W}{\Delta t}$
Units of Power	○ 1Watt = 1J/s = 1kgm²/s³