KE = ½ mv²

Question 1

The law of conversation of energy tell us that energy cannot be:

- A. changed in form.
- B. created or destroyed.
- C. measured.
- D. calculated.

Question 1

The law of conversation of energy tell us that energy cannot be:

- A. changed in form.
- B. created or destroyed.
- C. measured.
- D. calculated.

Question 2

Under what conditions would a 50kg cannonball have the most kinetic energy?

- A. The cannonball is at rest at ground level
- B. The cannonball has a speed of 10 m/s and is at ground level.
- C. The cannonball is at rest 10 meters above the ground.
- D. The cannonball has a speed of 5 m/s and is 5 meters above the

Question 2

Under what conditions would a 50kg cannonball have the most kinetic energy?

- A. The cannonball is at rest at ground level
- B. The cannonball has a speed of 10 m/s and is at ground level.
- C. The cannonball is at rest 10 meters above the ground.
- D. The cannonball has a speed of 5 m/s and is 5 meters above the ground.

Question 3

- A. The cannonball is at rest at ground level
- B. The cannonball has a speed of 10 m/s and is at ground level.
- C. The cannonball is at rest 10 meters above the ground.
- D. The cannonball has a speed of 5 m/s and is 5 meters above the ground.

Question 3


Under what conditions would a 50kg cannonball have the most potential energy?

- A. The cannonball is at rest at ground level
- B. The cannonball has a speed of 10 m/s and is at ground level.
- C. The cannonball is at rest 10 meters above the ground.
- D. The cannonball has a speed of 5 m/s and is 5 meters above the ground.

GPE = mgh

Grade 8 Only

Question 4




A 20 kg mass is attached a spring at rest as in the image above. The spring is then stretched 1 m to the right, and is released. At the instant the mass has moved 0.5 m back, what types of mechanical energy does $\,$ the mass possess?

- A. Gravitational potential only.
- B. Kinetic only
- C. Spring potential only.
- D. Kinetic and spring potential only

Grade 8 Only

Question 4



A 20 kg mass is attached a spring at rest as in the image above. The spring is then stretched 1 m to the right, and is released. At the instant the mass has moved 0.5 m back, what types of mechanical energy does the mass possess?

- A. Gravitational potential only.
- B. Kinetic only
- C. Spring potential only.
- D. Kinetic and spring potential only

Obviously still moving and not yet back to its equilibrium position so still has SPE.

Question 5

A box of old textbooks is on the middle shelf in the bookroom 1.3 m from the floor. If the janitor relocates the box to a shelf that is 2.6 \mbox{m} from the floor, how much work does he do on the box? m = 10 kg.

- A. 13 J
- B. 26 J
- C. 130 J
- D. 60 J

Question 5

A box of old textbooks is on the middle shelf in the bookroom 1.3 m from the floor. If the janitor relocates the box to a shelf that is 2.6 m from the floor, how much work does he do on the box? m = 10 kg.

- A. 13 J
- GPE = mgh = $10kg * 10m/s^2 * 1.3m$
- B. 26 J C. 130 J
- D. 60 J

Momentum so Grade 8 Only for now

Question 6

A skateboard moves along a frictionless level road at a constant speed. A kid falls straight down onto it. What happens to the velocity of the skateboard?

- A. It increases
- B. It remains the same
- C. It decreases
- D. it stops immediately

Momentum so Grade 8 Only for now

Question 6

A skateboard moves along a frictionless level road at a constant speed. A kid falls straight down onto it. What happens to the velocity of the skateboard?

A. It increases

Conservation of momentum

B. It remains the same C. It decreases

 $m_s v_i = (m_s + m_k) v_f$

D. it stops immediately

As total mass has increased so $v_f < v_i$ to keep the

momentum the same.

Question 7

If you push for a half hour or a whole hour against a stationary wall

- A. twice as much work is done during the half hour.
- B. no work on the wall is done in either case.
- C. half as much work is done during the half hour.
- D. it is impossible to determine how much work is done.

Question 7

If you push for a half hour or a whole hour against a stationary wall

- A. twice as much work is done during the half hour.
- B. no work on the wall is done in either case.

W = Fd

- C. half as much work is done during the half hour.
- D. it is impossible to determine how much work is done.

Question 8

If you push an object twice as far while applying the same force, you do

- A. four times as much work.
- B. twice as much work.
- C. the same amount of work.
- D. half as much work.

Question 8

If you push an object twice as far while applying the same force, you do

- A. four times as much work.
- B. twice as much work.

W = Fd

- C. the same amount of work.
- D. half as much work.

Grade 8 Only

Question 9

A bow is drawn so that it has 50 J of potential energy stored in the bow. Neglecting air resistance, when the bow is fired, the arrow will have a kinetic energy that is

- A. less than 50 J.
- B. 50 J.
- C. more than 50 J.
- D. impossible to predict without additional information

Grade 8 Only

Question 9

A bow is drawn so that it has 50 J of potential energy stored in the bow. Neglecting air resistance, when the bow is fired, the arrow will have a kinetic energy that is

A. less than 50 J.

Conservation of Mechanical Energy

B. 50 J.

ME_i = ME_t

SPE_i = KE_t (initially no KE, after firing no SPE)

C. more than 50 J.

D. impossible to predict without additional information

Question 10

A skier skiing down a frictionless incline has its maximum speed

- A. halfway down.
- B. at the top.
- C. at the bottom.
- D. depends upon the incline of the ramp.

Question 10

A skier skiing down a frictionless incline has its maximum speed

- A. halfway down.
- B. at the top. C. at the bottom.

Conservation of Mechanical Energy

 $\mathrm{ME_i} = \mathrm{ME_f}$ $\mathrm{GPE_i} = \mathrm{KE_f}$ (initially no KE, at the bottom no GPE)

D. depends upon the incline of the ramp.

Question 11

It takes 40 J to push a large box 4 m across a floor. Assuming the push is in the same direction as the move, what is the magnitude of the force on the box?

- A. 10 N
- B. 40 N
- C. 160 N
- D. 4 N

Question 11

It takes 40 J to push a large box 4 m across a floor. Assuming the push is in the same direction as the move, what is the magnitude of the force on the box?

A. 10 N

W = Fd F = W/d = 40J/4m =

B. 40 N C. 160 N

D. 4 N

Question 12

A truck moves 4 times as fast as another identical truck. Compared to the slower truck, the faster truck has

- A. 8 times the KE.
- B. 16 times the KE.
- C. 12 times the KE.
- D. 4 times the KE.

Question 12

A truck moves 4 times as fast as another identical truck. Compared to the slower truck, the faster truck has

- A. 8 times the KE.
- B. 16 times the KE.

KE = ½ mv²

- C. 12 times the KE.
- D. 4 times the KE.

Grade 8 Only

Question 13

Spring A is extended 4 times as far as another identical spring, called spring B. Compared to the spring B, spring A has

- A. 8 times the PEspring.
- B. 16 times the PEspring.
- C. 12 times the PEspring.
- D. 4 times the PEspring.

Grade 8 Only

Question 13

Spring A is extended 4 times as far as another identical spring, called spring B. Compared to the spring B, spring A has

- A. 8 times the PEspring.
- A. 8 times the PEspring. B. 16 times the PEspring. $E_p = \frac{1}{2}kx^2$ or $PE_{elastic} = \frac{1}{2}kx^2$
- C. 12 times the PEspring.
- D. 4 times the PEspring.

Question 14

Which has greater kinetic energy, a car traveling at 30 km/hr or a car of half the mass traveling at 60 km/hr?

- A. the 30 km/hr car
- B. the 60 km/hr car
- C. Both have the same kinetic energy.
- D. More information is needed about the distance traveled.

Question 14

Which has greater kinetic energy, a car traveling at 30 km/hr or a car of half the mass traveling at 60 km/hr?

- A. the 30 km/hr car
- B. the 60 km/hr car
- $KE = \frac{1}{2} mv^2$
- C. Both have the same kinetic energy.
- D. More information is needed about the distance traveled.

Question 15

A diver who has a mass of 50 kg steps off a diving board that is 10 m $\,$ above the water. The diver hits the water with kinetic energy of

- A. 500 J.
- B. 510 J.
- C. 5000 J.
- D. 10 J.

Question 15

A diver who has a mass of 50 kg steps off a diving board that is 10 m above the water. The diver hits the water with kinetic energy of

- A. 500 J.
- B. 510 J.

Conservation of Mechanical Energy

- C. 5000 J.
- D. 10 J.

 $\mathrm{ME_{i}} = \mathrm{ME_{f}}$ GPE $_{i} = \mathrm{KE_{f}}$ (initially no KE, when hits the water no GPE)

Question 16

Which of the following represents the units for energy in base SI units?

B.
$$\frac{kgn}{2}$$

C.
$$\frac{kgm^2}{s}$$

D.
$$\frac{kgm^2}{s^2}$$

Question 16

Which of the following represents the units for energy in base SI units?

$$\frac{kgm}{}$$
 F = ma = kgm/s²

B.
$$\frac{kgm}{s^2}$$
 W = Fd = kgm/s² . m = kgm²/s²

C.
$$\frac{kgm^2}{s}$$
 KE = ½ mv² = kgm/s . m/s = kgm²/s²

$$S^2$$
 GPE = mgh = kg · m/s² · m = kgm²/s²

Question 17

Which of the following is the energy stored in an object by virtue of its position?

- A. Potential energy
- B. Gravitational potential energy
- C. Spring potential energy
- D. Kinetic energy

Question 17

Which of the following is the energy stored in an object by virtue of its position?

- A. Potential energy
- B. Gravitational potential energy
- C. Spring potential energy
- D. Kinetic energy

Grade 8 Only

Question 18

A force of 10 N stretches a spring that has a spring constant of 20 N/m. The spring obeys Hooke's law.

What is the extension of the spring?

- A. 0.5 m
- B. 200 m
- C. 2 m
- D. 20 m

Grade 8 Only

Question 18

A force of 10 N stretches a spring that has a spring constant of 20 N/m. The spring obeys Hooke's law.

What is the extension of the spring?

- A. 0.5 m
- F = kx x = F/k
- B. 200 m C. 2 m
- = 10N / 20 N/m
- D. 20 m

Grade 8 Only

Question 19

A force of 10 N stretches a spring that has a spring constant of 20 N/m. The spring obeys Hooke's law.

What is the potential energy stored in the spring?

A. 2.5 J

B. 5.0 J

C. 10 J

D. 200 J

Grade 8 Only

Question 19

A force of 10 N stretches a spring that has a spring constant of 20 N/m. The spring obeys Hooke's law.

What is the potential energy stored in the spring?

A. 2.5 J F = kx

B. 5.0 J x = F/k= 10N / 20 N/m

C. 10 J

D. 200 J $E_{spring} = \frac{1}{2} kx^2 = \frac{1}{2} * 20 * 0.5^2$